

For More Information

Articles

Adubofour K, Keenan CR, Daftary A, Mensah-Adubofour J, Dachman WD. Strategies to reduce medication errors in ambulatory practice. *J Natl Med Assoc.* 2004;96:1558-1564.

<http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2568660/?page=1>

Bosk CL, Dixon-Woods M, Goeschel CA, Pronovost PJ. Reality check for checklists. *Lancet.* 2009;374:444-445.

[http://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(09\)61440-9/fulltext](http://www.thelancet.com/journals/lancet/article/PIIS0140-6736(09)61440-9/fulltext)

Budnitz DA, Pollock DA, Weidenback KN, et al. National surveillance of emergency department visits for outpatient adverse drug events. *JAMA.* 2006;296:1858-1866.

<http://jama.ama-assn.org/cgi/content/full/296/15/1858?max-toshow=&HITS=10&hits=10&RESULTFORMAT=&fulltext=National+Surveillance+of+Emergency+Department+Visits+for+Outpatient+Adverse+Drug+Events%2C+&searchid=1&FIRSTINDEX=0&resourcetype=HWCIT>

Casalino LP, Dunham D, Chin MH, et al. Frequency of failure to inform patients of clinically significant outpatient test results. *Arch Intern Med.* 2009;169:1123-1129.

http://archinte.amaassn.org/cgi/search?fulltext=Frequency+of+failure+to+inform+patients+of+clinically+significant+outpatient+test+results.+&quicksearch_submit.x=7&quicksearch_submit.y=6

Dovey SM, Meyers DS, Phillips RL Jr, et al. A preliminary taxonomy of medical errors in family practice. *Qual Saf Health Care.* 2002;11:233-238.

<http://qshc.bmj.com/content/11/3/233.abstract>

Forster AJ, Murff HJ, Peterson JF, Gandhi TK, Bates DW. The incidence and severity of adverse events affecting patients after discharge from the hospital. *Ann Intern Med.* 2003;138:161-167. <http://www.annals.org/content/138/3/161.full.pdf+html>

Levinson W, Roter DL, Mullooly JP, Dull VT, Frankel RM. Physician-patient communication: the relationship with malpractice claims among primary care physicians and surgeons. *JAMA.* 1997;277:553-559. <http://jama.ama-assn.org/cgi/reprint/277/7/553>

Newman-Toker DE, Pronovost PJ. Diagnostic errors—the next frontier for patient safety. *JAMA.* 2009; 301:1060-1062. <http://jama.ama-assn.org/cgi/reprint/301/10/1060>

Phillips RL Jr, Bartholomew LA, Dovey SM, et al. Learning from malpractice claims about negligent, adverse events in primary care in the United States. *Qual Saf Health Care.* 2004;13:121-126.

Schiff GD, Hasan O, Seijeoung K, et al. Diagnostic error in medicine: analysis of 583 physician-reported errors. *Arch Intern Med.* 2009;169:1881-1887. <http://archinte.ama-assn.org/cgi/content/full/169/20/1881>

Smith PC, Araya-Guerra R, Bublitz C, et al. Missing clinical information during primary care visits. *JAMA.* 2005;293:565-571. <http://jama.ama-assn.org/cgi/content/full/293/5/565>

Solberg LI, Elward KS, Phillips WR, et al. How can primary care cross the quality chasm? *Ann Fam Med.* 2009;7:164-169. <http://www.annfammed.org/cgi/content/full/7/2/164>

Thomsen LA, Winterstein AG, Søndergaard B, Haugbølle LS, Melander A. Systematic review of the incidence and characteristics of preventable adverse drug events in ambulatory care. *Ann Pharmacother.* 2007;41:1411-1426. <http://www.theannals.com/cgi/content/abstract/41/9/1411>

Websites

The Agency for Healthcare Research and Quality's home page includes links to online journals and newsletters, tools for improving patient safety, and notices of conferences and workshops. There is also useful information at the agency's Patient Safety Net home page and on its Medical Office Patient Safety Culture Survey page.

Home page: <http://www.ahrq.gov/>

Patient Safety Net: <http://psnet.ahrq.gov/>

Patient Safety Culture Survey:
<http://www.ahrq.gov/qual/mosurvey08/>

The Centers for Medicare and Medicaid Services provides information about e-prescribing and health information technology at the following sites:

<http://www.cms.hhs.gov/EPrescribing> and

<http://healthit.hhs.gov/portal/server.pt>

The Certification Commission for Health Information Technology has a list of EHR products that have earned what's likely to be the federal seal of approval for "certification" under the HITECH Act.

<http://www.cchit.org>

The Electronic Prescribing Readiness Assessment helps you determine whether your practice is ready for e-prescribing.

<http://www.getrxconnected.com/>

Sources for sample policy-and-procedure manuals

The following professional associations provide background on EHRs and other error-proofing resources:

■ American Academy of Family Physicians

<http://www.aafp.org/online/en/home/practicemgt.html>

■ American Academy of Pediatrics

<http://practice.aap.org/>

■ American College of Physicians

http://www.acponline.org/running_practice/practice_management/

Brief Summary: For complete details, please see full Prescribing Information for Lexapro.

WARNINGS: SUICIDALITY AND ANTIDEPRESSANT DRUGS

Antidepressants increased the risk compared to placebo of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults in short-term studies of major depressive disorder (MDD) and other psychiatric disorders. Anyone considering the use of Lexapro or any other antidepressant in a child, adolescent, or young adult must balance this risk with the clinical need. Short-term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24; there was a reduction in risk with antidepressants compared to placebo in adults aged 65 and older. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide. Patients of all ages who are started on antidepressant therapy should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior. Families and caregivers should be advised of the need for close observation and communication with the prescriber. Lexapro is not approved for use in pediatric patients less than 12 years of age. (See Warnings and Precautions: Clinical Worsening and Suicide Risk, Patient Counseling Information: Information for Patients, and Use in Specific Populations: Pediatric Use).

INDICATIONS AND USAGE: Major Depressive Disorder—Lexapro (escitalopram) is indicated for the acute and maintenance treatment of major depressive disorder in adults and in adolescents 12 to 17 years of age (see *Clinical Studies*). A major depressive episode (DSM-IV) implies a prominent and relatively persistent (nearly every day for at least 2 weeks) depressed or dysphoric mood that usually interferes with daily functioning, and includes at least five of the following nine symptoms: depressed mood, loss of interest in usual activities, significant change in weight and/or appetite, insomnia or hypersomnia, psychomotor agitation or retardation, increased fatigue, feelings of guilt or worthlessness, slowed thinking or impaired concentration, a suicide attempt or suicidal ideation. **Generalized Anxiety Disorder—Lexapro** is indicated for the acute treatment of Generalized Anxiety Disorder (GAD) in adults (see *Clinical Studies*). Generalized Anxiety Disorder (DSM-IV) is characterized by excessive anxiety and worry (apprehensive expectation) that is persistent for at least 6 months and which the person finds difficult to control. It must be associated with at least 3 of the following symptoms: restlessness or feeling keyed up or on edge, being easily fatigued, difficulty concentrating or mind going blank, irritability, muscle tension, and sleep disturbance.

CONTRAINDICATIONS: Monoamine oxidase inhibitors (MAOIs)—Concomitant use in patients taking monoamine oxidase inhibitors (MAOIs) is contraindicated (see *Warnings and Precautions*). **Pimozide—Concomitant use in patients taking pimozide is contraindicated** (see *Drug Interactions*). **Hypersensitivity to escitalopram or citalopram—Lexapro is contraindicated in patients with a hypersensitivity to escitalopram or citalopram or any of the inactive ingredients in Lexapro.**

WARNINGS AND PRECAUTIONS: Clinical Worsening and Suicide Risk—Patients with major depressive disorder (MDD), both adult and pediatric, may experience worsening of their depression and/or the emergence of suicidal ideation and behavior (suicidality) or unusual changes in behavior, whether or not they are taking antidepressant medications, and this risk may persist until significant remission occurs. Suicide is a known risk of depression and certain other psychiatric disorders, and these disorders themselves are the strongest predictors of suicide. There has been a long-standing concern, however, that antidepressants may have a role in inducing worsening of depression and the emergence of suicidality in certain patients during the early phases of treatment. Pooled analyses of short-term placebo-controlled trials of antidepressant drugs (SSRIs and others) showed that these drugs increase the risk of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults (ages 18-24) with major depressive disorder (MDD) and other psychiatric disorders. Short-term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24; there was a reduction with antidepressants compared to placebo in adults aged 65 and older. The pooled analyses of placebo-controlled trials in children and adolescents with MDD, obsessive compulsive disorder (OCD), or other psychiatric disorders included a total of 24 short-term trials of antidepressant drugs in over 4400 patients. The pooled analyses of placebo-controlled trials in adults with MDD or other psychiatric disorders included a total of 295 short-term trials (median duration of 2 months) of 11 antidepressant drugs in over 77,000 patients. There was considerable variation in risk of suicidality among drugs, but a tendency toward an increase in the younger patients for almost all drugs studied. There were differences in absolute risk of suicidality across the different indications, with the highest increase in MDD. The risk differences (drug vs. placebo), however, were relatively stable within age strata and across indications. These risk differences (drug-placebo difference in the number of cases of suicidality per 1000 patients treated) are provided in Table 1.

TABLE 1	
Age Range	Drug-Placebo Difference in Number of Cases of Suicidality per 1000 Patients Treated
	Increases Compared to Placebo
<18	14 additional cases
18-24	5 additional cases
	Decreases Compared to Placebo
25-64	1 fewer case
≥65	6 fewer cases

No suicides occurred in any of the pediatric trials. There were suicides in the adult trials, but the number was not sufficient to reach any conclusion about drug effect on suicide. It is unknown whether the suicidality risk extends to longer-term use, i.e., beyond several months. However, there is substantial evidence from placebo-controlled maintenance trials in adults with depression that the use of antidepressants can delay the recurrence of depression. All patients being treated with antidepressants for any indication should be monitored appropriately and observed closely for clinical worsening, suicidality, and unusual changes in behavior, especially during the initial few months of a course of drug therapy, or at times of dose changes, either increases or decreases. The following symptoms, anxiety, agitation, panic attacks, insomnia, irritability, hostility, aggressiveness, impulsivity, akathisia (psychomotor restlessness), hypomania, and mania, have been reported in adult and pediatric patients being treated with antidepressants for major depressive disorder as well as for other indications, both psychiatric and nonpsychiatric. Although a causal link between the emergence of such symptoms and either the worsening of depression and/or the emergence of suicidal impulses has not been established, there is concern that such symptoms may represent precursors to emerging suicidality. Consideration should be given to changing the therapeutic regimen, including possibly discontinuing the medication, in patients whose depression is persistently worse, or who are experiencing emergent suicidality or symptoms that might be precursors to worsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's presenting symptoms. If the decision has been made to discontinue treatment, medication should be tapered, as rapidly as is feasible, but with recognition that abrupt discontinuation can be associated with certain symptoms (see *Dosage and Administration*). Families and caregivers of patients being treated with antidepressants for major depressive disorder or other indications, both psychiatric and nonpsychiatric, should be alerted about the need to monitor patients for the emergence of agitation, irritability, unusual changes in behavior, and the other symptoms described above, as well as the emergence of suicidality, and to report such symptoms immediately to health care providers. Such monitoring should include daily observation by families and caregivers (see also *Patient Counseling Information*). Prescriptions for Lexapro should be written for the smallest quantity of tablets consistent with good patient management, in order to reduce the risk of overdose. **Screening Patients for Bipolar Disorder—A major depressive episode may be the initial presentation of bipolar disorder. It is generally believed (though not established in controlled trials) that treating such an episode with an antidepressant alone may increase the likelihood of precipitation of a mixed/manic episode in patients at risk for bipolar disorder. Whether any of the symptoms described above represent such a conversion is unknown. However, prior to initiating treatment with an antidepressant, patients with depressive symptoms should be adequately screened to determine if they are at risk for bipolar disorder; such screening should include a detailed psychiatric history, including a family history of suicide, bipolar disorder, and depression. It should be noted that Lexapro is not approved for use in treating bipolar depression.**

Serotonin Syndrome or Neuroleptic Malignant Syndrome (NMS)-like Reactions—The development of a potentially life-threatening serotonin syndrome or Neuroleptic Malignant Syndrome (NMS)-like reactions have been reported with SSRIs and SSRI alone, including Lexapro treatment, but particularly with concomitant use of serotonergic drugs (including triptans) with drugs which impair metabolism of serotonin (including MAOIs), with antipsychotics or other dopamine antagonists. Serotonin syndrome symptoms may include mental status changes (e.g., agitation, hallucinations, coma), autonomic instability (e.g., tachycardia, labile blood pressure, hyperthermia), neuromuscular aberrations (e.g., hyperreflexia, incoordination) and/or gastrointestinal symptoms (e.g., nausea, vomiting, diarrhea). Serotonin syndrome, in its most severe form can resemble neuroleptic malignant syndrome, which includes hyperthermia, muscle rigidity, autonomic instability with possible rapid fluctuation of vital signs, and mental status changes. Patients should be monitored for the emergence of serotonin syndrome or NMS-like signs and symptoms. The concomitant use of Lexapro with MAOIs intended to treat depression is contraindicated. If concomitant treatment of Lexapro with a 5-hydroxytryptamine receptor agonist (triptan) is clinically warranted, careful observation of the patient is advised, particularly during treatment initiation and dose increases. The concomitant use of Lexapro with serotonin precursors (such as tryptophan) is not recommended. Treatment with Lexapro and any concomitant serotonergic or antidepressant agents, including antipsychotics, should be discontinued immediately if the above events occur and supportive symptomatic treatment should be initiated. **Discontinuation of Treatment with Lexapro—**During marketing of Lexapro and other SSRIs and SNRIs (serotonin and norepinephrine reuptake inhibitors), there have been spontaneous reports of serious events occurring upon discontinuation of these drugs, particularly when abrupt, including the following: dysphoric mood, irritability, agitation, dizziness, sensory disturbances (e.g., paresthesias such as electric shock sensations), anxiety, confusion, headache, lethargy, emotional lability, insomnia, and hypomania. While these events are generally self-limiting, there have been reports of serious discontinuation symptoms. Patients should be monitored for these symptoms when discontinuing treatment with Lexapro. A gradual reduction in the dose rather than abrupt cessation is recommended whenever possible. If intolerable symptoms occur following a decrease in the dose or upon discontinuation of treatment, then resuming the previously prescribed dose may be considered. Subsequently, the physician may continue decreasing the dose but at a more gradual rate (see *Dosage and Administration*). **Seizures—**Although anti-convulsant effects of racemic citalopram have been observed in animal studies, Lexapro has not been systematically evaluated in patients with a seizure disorder. These patients were excluded from clinical studies during the product's premarketing testing. In clinical trials of Lexapro, cases of convulsion have been reported in association with Lexapro treatment. Like other drugs effective in the treatment of major depressive disorder, Lexapro should be introduced with care in patients with a history of seizure disorder.

Activation of Mania/Hypomania—In placebo-controlled trials of Lexapro in major depressive disorder, activation of mania/hypomania was reported in one (0.1%) of 715 patients treated with Lexapro and in none of the 592 patients treated with placebo. One additional case of hypomania has been reported in association with Lexapro treatment. Activation of mania/hypomania has also been reported in a small proportion of patients with major affective disorders treated with racemic citalopram and other marketed drugs effective in the treatment of major depressive disorder. As with all drugs effective in the treatment of major depressive disorder, Lexapro should be used cautiously in patients with a history of mania. **Hypонатremia—**Hypонатremia may occur as a result of treatment with SSRIs and SNRIs, including Lexapro. In many cases, this hypонатremia appears to be the result of the syndrome of inappropriate antidiuretic hormone secretion (SIADH), and was reversible when Lexapro was discontinued. Cases with serum sodium lower than 110 mmol/L have been reported. Elderly patients may be at greater risk of developing hypонатremia with SSRIs and SNRIs. Also, patients taking diuretics or who are otherwise volume depleted may be at greater risk (see *Geriatric Use*). Discontinuation of Lexapro should be considered in patients with symptomatic hypонатremia and appropriate medical intervention should be instituted. Signs and symptoms of hypонатremia include headache, difficulty concentrating, memory impairment, confusion, weakness, and unsteadiness, which may lead to falls. Signs and symptoms associated with more severe and/or acute cases have included hallucination, syncope, seizure, coma, respiratory arrest, and death. **Abnormal Bleeding—**SSRIs and SNRIs, including Lexapro, may increase the risk of bleeding events. Concomitant use of aspirin, nonsteroidal anti-inflammatory drugs, warfarin, and other anticoagulants may add to the risk. Case reports and epidemiological studies (case-control and cohort design) have demonstrated an association between use of drugs that interfere with serotonin reuptake and the occurrence of gastrointestinal bleeding. Bleeding events related to SSRIs and SNRIs use have ranged from ecchymoses, hematomas, epistaxis, and petechiae to life-threatening hemorrhages. Patients should be cautioned about the risk of bleeding associated with the concomitant use of Lexapro and NSAIDs, aspirin, or other drugs that affect coagulation. **Interference with Cognitive and Motor Performance—**In a study

LEXAPRO® (escitalopram oxalate) TABLETS/ORAL SOLUTION

in normal volunteers, Lexapro 10 mg/day did not produce impairment of intellectual function or psychomotor performance. Because any psychoactive drug may impair judgment, thinking, or motor skills, however, patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that Lexapro therapy does not affect their ability to engage in such activities. **Use in Patients with Concomitant Illness**-Clinical experience with Lexapro in patients with certain concomitant systemic illnesses is limited. Caution is advisable in using Lexapro in patients with diseases or conditions that produce altered metabolism or hemodynamic responses. Lexapro has not been systematically evaluated in patients with a recent history of myocardial infarction or unstable heart disease. Patients with these diagnoses were generally excluded from clinical studies during the product's premarketing testing. In subjects with hepatic impairment, clearance of racemic citalopram was decreased and plasma concentrations were increased. The recommended dose of Lexapro in hepatically impaired patients is 10 mg/day [see *Dosage and Administration*]. Because escitalopram is extensively metabolized, excretion of unchanged drug in urine is a minor route of elimination. Until adequate numbers of patients with severe renal impairment have been evaluated during chronic treatment with Lexapro, however, it should be used with caution in such patients [see *Dosage and Administration*]. **Potential for Interaction with Monoamine Oxidase Inhibitors**-In patients receiving serotonergic reuptake inhibitor drugs in combination with a monoamine oxidase inhibitor (MAOI), there have been reports of serious, sometimes fatal, reactions including hyperthermia, rigidity, myoclonus, autonomic instability with possible rapid fluctuations of vital signs, and mental status changes that include extreme agitation progressing to delirium and coma. These reactions have also been reported in patients who have recently discontinued SSRI treatment and have been started on a MAOI. Some cases presented with features resembling neuroleptic malignant syndrome. Furthermore, limited animal data on the effects of combined use of SSRIs and MAOIs suggest that these drugs may act synergistically to elevate blood pressure and evoke behavioral excitation. Therefore, it is recommended that Lexapro should not be used in combination with a MAOI, or within 14 days of discontinuing treatment with a MAOI. Similarly, at least 14 days should be allowed after stopping Lexapro before starting an MAOI. Serotonin syndrome has been reported in two patients who were concomitantly receiving linezolid, an antibiotic which is a reversible non-selective MAOI.

ADVERSE REACTIONS: Clinical Trials Experience-Because clinical studies are conducted under widely varying conditions, adverse reaction rates observed in the clinical studies of a drug cannot be directly compared to rates in the clinical studies of another drug and may not reflect the rates observed in practice. **Clinical Trial Data Sources; Pediatrics (6-17 years)**-Adverse events were collected in 576 pediatric patients (286 Lexapro, 290 placebo) with major depressive disorder in double-blind placebo-controlled studies. Safety and effectiveness of Lexapro in pediatric patients less than 12 years of age has not been established. **Adults**-Adverse event information for Lexapro was collected from 715 patients with major depressive disorder who were exposed to escitalopram and from 592 patients who were exposed to placebo in double-blind, placebo-controlled trials. An additional 284 patients with major depressive disorder were newly exposed to escitalopram in open-label trials. The adverse event information for Lexapro in patients with GAD was collected from 429 patients exposed to escitalopram and from 427 patients exposed to placebo in double-blind, placebo-controlled trials. Adverse events during exposure were obtained primarily by general inquiry and recorded by clinical investigators using terminology of their own choosing. Consequently, it is not possible to provide a meaningful estimate of the proportion of individuals experiencing adverse events without first grouping similar types of events into a smaller number of standardized event categories. In the tables and tabulations that follow, standard World Health Organization (WHO) terminology has been used to classify reported adverse events. The stated frequencies of adverse reactions represent the proportion of individuals who experienced, at least once, a treatment-emergent adverse event of the type listed. An event was considered treatment-emergent if it occurred for the first time or worsened while receiving therapy following baseline evaluation. **Adverse Events Associated with Discontinuation of Treatment; Major Depressive Disorder; Pediatrics (6-17 years)**-Adverse events were associated with discontinuation of 3.5% of 286 patients receiving Lexapro and 1% of 290 patients receiving placebo. The most common adverse event (incidence at least 1% for Lexapro and greater than placebo) associated with discontinuation was insomnia (1% Lexapro, 0% placebo). **Adults**-Among the 715 depressed patients who received Lexapro in placebo-controlled trials, 6% discontinued treatment due to an adverse event, as compared to 2% of 592 patients receiving placebo. In two fixed-dose studies, the rate of discontinuation for adverse events in patients receiving 10 mg/day Lexapro was not significantly different from the rate of discontinuation for adverse events in patients receiving placebo. The rate of discontinuation for adverse events in patients assigned to a fixed dose of 20 mg/day Lexapro was 10%, which was significantly different from the rate of discontinuation for adverse events in patients receiving 10 mg/day Lexapro (4% and placebo (3%). Adverse events that were associated with the discontinuation of at least 1% of patients treated with Lexapro, and for which the rate was at least twice that of placebo, were nausea (2%) and ejaculation disorder (2% of male patients). **Generalized Anxiety Disorder; Adults**-Among the 429 GAD patients who received Lexapro 10-20 mg/day in placebo-controlled trials, 8% discontinued treatment due to an adverse event, as compared to 4% of 427 patients receiving placebo. Adverse events that were associated with the discontinuation of at least 1% of patients treated with Lexapro, and for which the rate was at least twice the placebo rate, were nausea (2%), insomnia (1%), and fatigue (1%). **Incidence of Adverse Reactions in Placebo-Controlled Clinical Trials; Major Depressive Disorder; Pediatrics (6-17 years)**-The overall profile of adverse reactions in pediatric patients was generally similar to that seen in adult studies, as shown in Table 2. However, the following adverse reactions (excluding those which appear in Table 2 and those for which the coded terms were uninformative or misleading) were reported at an incidence of at least 2% for Lexapro and greater than placebo: back pain, urinary tract infection, vomiting, and nasal congestion. **Adults**-The most commonly observed adverse reactions in Lexapro patients (incidence of approximately 5% or greater and approximately twice the incidence in placebo patients) were insomnia, ejaculation disorder (primarily ejaculatory delay), nausea, sweating increased, fatigue, and somnolence. Table 2 enumerates the incidence, rounded to the nearest percent, of treatment-emergent adverse events that occurred among 715 depressed patients who received Lexapro at doses ranging from 10 to 20 mg/day in placebo-controlled trials. Events included are those occurring in 2% or more of patients treated with Lexapro and for which the incidence in patients treated with Lexapro was greater than the incidence in placebo-treated patients.

TABLE 2
Treatment-Emergent Adverse Reactions Observed with a Frequency of $\geq 2\%$
and Greater Than Placebo for Major Depressive Disorder

Adverse Reaction	Lexapro (N=715)	Placebo (N=592)
Autonomic Nervous System Disorders		
Dry Mouth	6%	2%
Sweating Increased	5%	2%
Central & Peripheral Nervous System Disorders		
Dizziness	5%	3%
Gastrointestinal Disorders		
Nausea	15%	7%
Diarrhea	8%	5%
Constipation	3%	1%
Indigestion	3%	1%
Abdominal Pain	2%	1%
General		
Influenza-like Symptoms	5%	4%
Fatigue	5%	2%
Psychiatric Disorders		
Insomnia	9%	4%
Somnolence	6%	2%
Appetite Decreased	3%	1%
Libido Decreased	3%	1%
Respiratory System Disorders		
Rhinitis	5%	4%
Sinusitis	3%	2%
Urogenital		
Ejaculation Disorder ^{1,2}	9%	<1%
Impotence ²	3%	<1%
Anorgasmia ³	2%	<1%

¹Primarily ejaculatory delay.

²Denominator used was for males only (N=225 Lexapro; N=188 placebo).

³Denominator used was for females only (N=490 Lexapro; N=404 placebo).

Generalized Anxiety Disorder; Adults-The most commonly observed adverse reactions in Lexapro patients (incidence of approximately 5% or greater and approximately twice the incidence in placebo patients) were nausea, ejaculation disorder (primarily ejaculatory delay), insomnia, fatigue, decreased libido, and anorgasmia. Table 3 enumerates the incidence, rounded to the nearest percent of treatment-emergent adverse events that occurred among 429 GAD patients who received Lexapro 10 to 20 mg/day in placebo-controlled trials. Events included are those occurring in 2% or more of patients treated with Lexapro and for which the incidence in patients treated with Lexapro was greater than the incidence in placebo-treated patients.

LEXAPRO® (escitalopram oxalate) TABLETS/ORAL SOLUTION

TABLE 3

Treatment-Emergent Adverse Reactions Observed with a Frequency of $\geq 2\%$ and Greater Than Placebo for Generalized Anxiety Disorder

Adverse Reactions	Lexapro (N=429)	Placebo (N=427)
Autonomic Nervous System Disorders		
Dry Mouth	9%	5%
Sweating Increased	4%	1%
Central & Peripheral Nervous System Disorders		
Headache	24%	17%
Paresthesia	2%	1%
Gastrointestinal Disorders		
Nausea	18%	8%
Diarrhea	8%	6%
Constipation	5%	4%
Indigestion	3%	2%
Vomiting	3%	1%
Abdominal Pain	2%	1%
Flatulence	2%	1%
Toothache	2%	0%
General		
Fatigue	8%	2%
Influenza-like Symptoms	5%	4%
Musculoskeletal System Disorder		
Neck/Shoulder Pain	3%	1%
Psychiatric Disorders		
Somnolence	13%	7%
Insomnia	12%	6%
Libido Decreased	7%	2%
Dreaming Abnormal	3%	2%
Appetite Decreased	3%	1%
Lethargy	3%	1%
Respiratory System Disorders		
Yawning	2%	1%
Urogenital		
Ejaculation Disorder ^{1,2}	14%	2%
Anorgasmia ³	6%	<1%
Menstrual Disorder	2%	1%

¹Primarily ejaculatory delay.

²Denominator used was for males only (N=182 Lexapro; N=195 placebo).

³Denominator used was for females only (N=247 Lexapro; N=232 placebo).

Dose Dependency of Adverse Reactions—The potential dose dependency of common adverse reactions (defined as an incidence rate of $\geq 5\%$ in either the 10 mg or 20 mg Lexapro groups) was examined on the basis of the combined incidence of adverse events in two fixed-dose trials. The overall incidence rates of adverse events in 10 mg Lexapro-treated patients (66%) was similar to that of the placebo-treated patients (61%), while the incidence rate in 20 mg/day Lexapro-treated patients was greater (86%). Table 4 shows common adverse reactions that occurred in the 20 mg/day Lexapro group with an incidence that was approximately twice that of the 10 mg/day Lexapro group and approximately twice that of the placebo group.

TABLE 4

Incidence of Common Adverse Reactions in Patients with Major Depressive Disorder

Adverse Reaction	Placebo (N=311)	10 mg/day Lexapro (N=310)	20 mg/day Lexapro (N=125)
Insomnia	4%	7%	14%
Diarrhea	5%	6%	14%
Dry Mouth	3%	4%	9%
Somnolence	1%	4%	9%
Dizziness	2%	4%	7%
Sweating Increased	<1%	3%	8%
Constipation	1%	3%	6%
Fatigue	2%	2%	6%
Indigestion	1%	2%	6%

Male and Female Sexual Dysfunction with SSRIs—Although changes in sexual desire, sexual performance, and sexual satisfaction often occur as manifestations of a psychiatric disorder, they may also be a consequence of pharmacologic treatment. In particular, some evidence suggests that SSRIs can cause such untoward sexual experiences. Reliable estimates of the incidence and severity of untoward experiences involving sexual desire, performance, and satisfaction are difficult to obtain, however, in part because patients and physicians may be reluctant to discuss them. Accordingly, estimates of the incidence of untoward sexual experience and performance cited in product labeling are likely to underestimate their actual incidence.

TABLE 5

Incidence of Sexual Side Effects in Placebo-Controlled Clinical Trials

Adverse Event	Lexapro	Placebo
	In Males Only	
	(N=407)	(N=383)
Ejaculation Disorder (primarily ejaculatory delay)	12%	1%
Libido Decreased	6%	2%
Impotence	2%	<1%
	In Females Only	
	(N=737)	(N=636)
Libido Decreased	3%	1%
Anorgasmia	3%	<1%

There are no adequately designed studies examining sexual dysfunction with escitalopram treatment. Priapism has been reported with all SSRIs. While it is difficult to know the precise risk of sexual dysfunction associated with the use of SSRIs, physicians should routinely inquire about such possible side effects. **Vital Sign Changes**—Lexapro and placebo groups were compared with respect to (1) mean change from baseline in vital signs (pulse, systolic blood pressure, and diastolic blood pressure) and (2) the incidence of patients meeting criteria for potentially clinically significant changes from baseline in these variables. These analyses did not reveal any clinically important changes in vital signs associated with Lexapro treatment. In addition, a comparison of supine and standing vital sign measures in subjects receiving Lexapro indicated that Lexapro treatment is not associated with orthostatic changes. **Weight Changes**—Patients treated with Lexapro in controlled trials did not differ from placebo-treated patients with regard to clinically important change in body weight. **Laboratory Changes**—Lexapro and placebo groups were compared with respect to (1) mean change from baseline in various serum chemistry, hematology, and urinalysis variables, and (2) the incidence of patients meeting criteria for potentially clinically significant changes from baseline in these

LEXAPRO® (escitalopram oxalate) TABLETS/ORAL SOLUTION

mortality. In a retrospective, case-control study of 377 women whose infants were born with PPHN and 836 women whose infants were born healthy, the risk for developing PPHN was approximately six-fold higher for infants exposed to SSRIs after the 20th week of gestation compared to infants who had not been exposed to antidepressants during pregnancy. There is currently no corroborative evidence regarding the risk for PPHN following exposure to SSRIs in pregnancy; this is the first study that has investigated the potential risk. The study did not include enough cases with exposure to individual SSRIs to determine if all SSRIs posed similar levels of PPHN risk. When treating a pregnant woman with Lexapro during the third trimester, the physician should carefully consider both the potential risks and benefits of treatment [see *Dosage and Administration*]. Physicians should note that in a prospective longitudinal study of 201 women with a history of major depression who were euthymic at the beginning of pregnancy, women who discontinued antidepressant medication during pregnancy were more likely to experience a relapse of major depression than women who continued antidepressant medication. **Labor and Delivery**—The effect of Lexapro on labor and delivery in humans is unknown. **Nursing Mothers**—Escitalopram is excreted in human breast milk. Limited data from women taking 10-20 mg escitalopram showed that exclusively breast-fed infants receive approximately 3.9% of the maternal weight-adjusted dose of escitalopram and 1.7% of the maternal weight-adjusted dose of desmethylcitalopram. There were two reports of infants experiencing excessive somnolence, decreased feeding, and weight loss in association with breastfeeding from a racemic citalopram-treated mother; in one case, the infant was reported to recover completely upon discontinuation of racemic citalopram by its mother and, in the second case, no follow-up information was available. Caution should be exercised and breastfeeding infants should be observed for adverse reactions when Lexapro is administered to a nursing woman. **Pediatric Use**—Safety and effectiveness of Lexapro has not been established in pediatric patients (less than 12 years of age) with Major Depressive Disorder. Safety and effectiveness of Lexapro has been established in adolescents (12 to 17 years of age) for the treatment of major depressive disorder [see *Clinical Studies*]. Although maintenance efficacy in adolescent patients with Major Depressive Disorder has not been systematically evaluated, maintenance efficacy can be extrapolated from adult data along with comparisons of escitalopram pharmacokinetic parameters in adults and adolescent patients. Safety and effectiveness of Lexapro has not been established in pediatric patients less than 18 years of age with Generalized Anxiety Disorder. **Geriatric Use**—Approximately 6% of the 1144 patients receiving escitalopram in controlled trials of Lexapro in major depressive disorder and GAD were 60 years of age or older; elderly patients in these trials received daily doses of Lexapro between 10 and 20 mg. The number of elderly patients in these trials was insufficient to adequately assess for possible differential efficacy and safety measures on the basis of age. Nevertheless, greater sensitivity of some elderly individuals to effects of Lexapro cannot be ruled out. SSRIs and SNRIs, including Lexapro, have been associated with cases of clinically significant hyponatremia in elderly patients, who may be at greater risk for this adverse event [see *Hyponatremia*]. In two pharmacokinetic studies, escitalopram half-life was increased by approximately 50% in elderly subjects as compared to young subjects and C_{24h} was unchanged [see *Clinical Pharmacology*], 10 mg/day is the recommended dose for elderly patients [see *Dosage and Administration*]. Of 4422 patients in clinical studies of racemic citalopram, 1357 were 60 and over, 1034 were 65 and over, and 457 were 75 and over. No overall differences in safety or effectiveness were observed between these subjects and younger subjects, and other reported clinical experience has not identified differences in responses between the elderly and younger patients, but again, greater sensitivity of some elderly individuals cannot be ruled out.

DRUG ABUSE AND DEPENDENCE: Abuse and Dependence; Physical and Psychological Dependence—Animal studies suggest that the abuse liability of racemic citalopram is low. Lexapro has not been systematically studied in humans for its potential for abuse, tolerance, or physical dependence. The premarketing clinical experience with Lexapro did not reveal any drug-seeking behavior. However, these observations were not systematic and it is not possible to predict on the basis of this limited experience the extent to which a CNS-active drug will be misused, diverted, and/or abused once marketed. Consequently, physicians should carefully evaluate Lexapro patients for history of drug abuse and follow such patients closely, observing them for signs of misuse or abuse (e.g., development of tolerance, increments of dose, drug-seeking behavior).

OVERDOSAGE: Human Experience—In clinical trials of escitalopram, there were reports of escitalopram overdose, including overdoses of up to 600 mg, with no associated fatalities. During the postmarketing evaluation of escitalopram, Lexapro overdoses involving overdoses of over 1000 mg have been reported. As with other SSRIs, a fatal outcome in a patient who has taken an overdose of escitalopram has been rarely reported. Symptoms most often accompanying escitalopram overdose, alone or in combination with other drugs and/or alcohol, included convulsions, coma, dizziness, hypotension, insomnia, nausea, vomiting, sinus tachycardia, somnolence, and ECG changes (including QT prolongation and very rare cases of torsade de pointes). Acute renal failure has been very rarely reported accompanying overdose. **Management of Overdose**—Establish and maintain an airway to ensure adequate ventilation and oxygenation. Gastric evacuation by lavage and use of activated charcoal should be considered. Careful observation and cardiac and vital sign monitoring are recommended, along with general symptomatic and supportive care. Due to the large volume of distribution of escitalopram, forced diuresis, dialysis, hemoperfusion, and exchange transfusion are unlikely to be of benefit. There are no specific antidotes for Lexapro. In managing overdose, consider the possibility of multiple-drug involvement. The physician should consider contacting a poison control center for additional information on the treatment of any overdose.

Forest Pharmaceuticals, Inc.
Subsidiary of Forest Laboratories, Inc.
St. Louis, MO 63045 USA
Licensed from H. Lundbeck A/S

Treat your patients with the proven efficacy of LEXAPRO¹⁻³

Indicated to treat MDD and GAD in adults and also MDD in adolescents aged 12 to 17



Lexapro (escitalopram oxalate) is indicated for the acute and maintenance treatment of major depressive disorder (MDD) in adults and adolescents aged 12-17 years. Lexapro is also indicated for the acute treatment of generalized anxiety disorder (GAD) in adults.

IMPORTANT SAFETY INFORMATION

Warnings and Precautions

- All patients treated with antidepressants should be monitored appropriately and observed closely for clinical worsening, suicidality and unusual changes in behavior, especially within the first few months of treatment or when changing the dose. Consideration should be given to changing the therapeutic regimen, including discontinuing medication, in patients whose depression is persistently worse, who are experiencing emergent suicidality or symptoms that might be precursors to worsening depression or suicidality, especially if these symptoms are severe, abrupt in onset, or were not part of the patient's presenting symptoms. Families and caregivers of patients treated with antidepressants should be alerted about the need to monitor patients daily for the emergence of agitation, irritability, unusual changes in behavior, or the emergence of suicidality, and report such symptoms immediately. Prescriptions for Lexapro should be written for the smallest quantity of tablets, consistent with good patient management, in order to reduce the risk of overdose.
- A major depressive episode may be the initial presentation of bipolar disorder. In patients at risk for bipolar disorder, treating such an episode with an antidepressant alone may increase the likelihood of precipitating a mixed/manic episode. Prior to initiating treatment with an antidepressant, patients should be adequately screened to determine if they are at risk for bipolar disorder. Lexapro should be used cautiously in patients with a history of mania or seizure disorder. Lexapro is not approved for use in treating bipolar depression.
- The concomitant use of Lexapro with other SSRIs, SNRIs, triptans, tryptophan, antipsychotics or other dopamine antagonists is not recommended due to potential development of life-threatening serotonin syndrome or neuroleptic malignant syndrome (NMS)-like reactions. Reactions have been reported with SNRIs and SSRIs alone, including Lexapro, but particularly with drugs that impair metabolism of serotonin (including MAOIs). Management of these events should include immediate discontinuation of Lexapro and the concomitant agent and continued monitoring.
- Patients should be monitored for adverse reactions when discontinuing treatment with Lexapro. During marketing of Lexapro and other SSRIs and SNRIs, there have been spontaneous reports of adverse events occurring upon discontinuation, including dysphoric mood, irritability, agitation, dizziness, sensory disturbances (e.g., paresthesias), anxiety, confusion, headache, lethargy, emotional lability, insomnia and hypomania. A gradual dose reduction rather than abrupt cessation is recommended whenever possible.
- SSRIs and SNRIs have been associated with clinically significant hyponatremia. Elderly patients and patients taking diuretics or who are otherwise volume-depleted appear to be at a greater risk. Discontinuation of Lexapro should be considered in patients with symptomatic hyponatremia and appropriate medical intervention should be instituted.
- SSRIs (including Lexapro) and SNRIs may increase the risk of bleeding. Patients should be cautioned that concomitant use of aspirin, NSAIDs, warfarin or other anticoagulants may add to the risk.
- Patients should be cautioned about operating hazardous machinery, including automobiles, until they are reasonably certain that Lexapro does not affect their ability to engage in such activities.
- Lexapro should be used with caution in patients with severe renal impairment or with diseases or conditions that alter metabolism or hemodynamic responses. In subjects with hepatic impairment, clearance of racemic citalopram was decreased and plasma concentrations were increased. The recommended dose of Lexapro in hepatically impaired patients is 10 mg/day.
- For pregnant or nursing mothers, Lexapro should be used only if the potential benefit justifies the potential risk to the fetus or child.

Adverse Reactions

- In clinical trials of MDD, the most common adverse reactions in adults treated with Lexapro (approximately 5% or greater and at least twice the incidence of placebo) were nausea (15% vs 7%), insomnia (9% vs 4%), ejaculation disorder (9% vs <1%), fatigue (5% vs 2%), somnolence (6% vs 2%), and increased sweating (5% vs 2%). In pediatric patients, the overall profile of adverse reactions was similar to that seen in adults; however, the following additional adverse reactions were reported at an incidence of at least 2% for Lexapro and greater than placebo: back pain, urinary tract infection, vomiting, and nasal congestion.
- In clinical trials of GAD, the most common adverse reactions in adults treated with Lexapro (approximately 5% or greater and at least twice the incidence of placebo) were nausea (18% vs 8%), ejaculation disorder (14% vs 2%), insomnia (12% vs 6%), fatigue (8% vs 2%), decreased libido (7% vs 2%) and anorgasmia (6% vs <1%).

References: 1. LEXAPRO [package insert]. St. Louis, Mo: Forest Pharmaceuticals, Inc.; 2009. 2. Burke WJ, Gergel I, Bose A. Fixed-dose trial of the single isomer SSRI escitalopram in depressed outpatients. *J Clin Psychiatry*. 2002;63:331-336. 3. Davidson JRT, Bose A, Korotzer A, Zheng H. Escitalopram in the treatment of generalized anxiety disorder: double-blind, placebo controlled, flexible-dose study. *Depress Anxiety*. 2004;19:234-240. 4. Wade A, Lemming OM, Hedegaard KB. Escitalopram 10 mg/day is effective and well tolerated in a placebo-controlled study in depression in primary care. *Int Clin Psychopharmacol*. 2002;17:95-102. 5. SDI, April 2008. Depression and Anxiety Treatment Market Overview. Based on longitudinal analysis of US electronic retail pharmacy claims submitted for third-party reimbursement. 6. Data on file, Forest Laboratories, Inc.

Please see additional Important Safety Information on the adjacent page.

Please see accompanying brief summary of Prescribing Information for LEXAPRO.

 Forest Pharmaceuticals, Inc.
Pharmaceuticals • Therapeutics • Healthcare • Devices • Managed Care • Specialty Sales

© 2009 Forest Laboratories, Inc. Printed in U.S.A. 41-1015537dR3 12/09

Lexapro
escitalopram oxalate 

Treat the core symptoms of Major Depressive Disorder (MDD)
& Generalized Anxiety Disorder (GAD) in adults

Depressed Mood • Tension • Anxiety

**ALSO
FDA APPROVED
for MDD in adolescents
aged 12 to 17¹**

See the effect of LEXAPRO

Proven efficacy in MDD
and GAD in adults.¹⁻³

- Significantly higher rates of response and remission vs placebo in adults^{3,4}
- Significantly improved quality-of-life (QOL) scores vs placebo in adults^{2,3}
- Prescribed to over 18 million US patients⁵
- Widely available on health plan formularies without restrictions⁶

There is no generic available for LEXAPRO

WARNING: SUICIDALITY AND ANTIDEPRESSANT DRUGS

Antidepressants increased the risk compared to placebo of suicidal thinking and behavior (suicidality) in children, adolescents, and young adults in short-term studies of major depressive disorder (MDD) and other psychiatric disorders. Anyone considering the use of Lexapro or any other antidepressant in a child, adolescent or young adult must balance this risk with the clinical need. Short-term studies did not show an increase in the risk of suicidality with antidepressants compared to placebo in adults beyond age 24; there was a reduction in risk with antidepressants compared to placebo in adults aged 65 and older. Depression and certain other psychiatric disorders are themselves associated with increases in the risk of suicide. Patients of all ages who are started on antidepressant therapy should be monitored appropriately and observed closely for clinical worsening, suicidality, or unusual changes in behavior. Families and caregivers should be advised of the need for close observation and communication with the prescriber. Lexapro is not approved for use in pediatric patients less than 12 years of age.

Contraindications

- Lexapro is contraindicated in patients taking monoamine oxidase inhibitors (MAOIs). There have been reports of serious, sometimes fatal, reactions with some cases resembling neuroleptic malignant syndrome (NMS) and serotonin syndrome. Features may include hyperthermia, rigidity, myoclonus, autonomic instability with possible rapid fluctuations of vital signs, and mental status changes that include extreme agitation progressing to delirium and coma. These reactions have also been reported in patients who have recently discontinued SSRI treatment and have been started on an MAOI. Serotonin syndrome was reported for two patients who were concomitantly receiving linezolid, an antibiotic which has MAOI activity. Lexapro should not be used in combination with an MAOI or within 14 days of discontinuing an MAOI. MAOIs should not be initiated within 14 days of discontinuing Lexapro.
- Lexapro is contraindicated in patients taking pimozide or with hypersensitivity to escitalopram or citalopram.

Lexapro
escitalopram oxalate 

Please see additional Important Safety Information on adjacent page.

Visit the LEXAPRO website at www.lexapro.com